

# Analysis of regulation of pentose utilisation in *Aspergillus niger* reveals evolutionary adaptations in *Eurotiales*

E. Battaglia<sup>1</sup>, L. Visser<sup>1</sup>, A. Nijssen<sup>1</sup>, G.J. van Veluw<sup>1</sup>, H.A.B. Wösten<sup>1</sup> and R.P. de Vries<sup>1,2\*</sup>

<sup>1</sup>Microbiology, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands; <sup>2</sup>CBS-KNAW, Fungal Biodiversity Centre, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands.

\*Correspondence: Ronald P. de Vries, [r.devries@cbs.knaw.nl](mailto:r.devries@cbs.knaw.nl)

**Abstract:** Aspergilli are commonly found in soil and on decaying plant material. D-xylose and L-arabinose are highly abundant components of plant biomass. They are released from polysaccharides by fungi using a set of extracellular enzymes and subsequently converted intracellularly through the pentose catabolic pathway (PCP).

In this study, the L-arabinose responsive transcriptional activator (AraR) is identified in *Aspergillus niger* and was shown to control the L-arabinose catabolic pathway as well as expression of genes encoding extracellular L-arabinose releasing enzymes. AraR interacts with the D-xylose-responsive transcriptional activator XlnR in the regulation of the pentose catabolic pathway, but not with respect to release of L-arabinose and D-xylose.

AraR was only identified in the *Eurotiales*, more specifically in the family *Trichocomaceae* and appears to have originated from a gene duplication event (from XlnR) after this order or family split from the other filamentous ascomycetes. XlnR is present in all filamentous ascomycetes with the exception of members of the *Onygenales*. Since the *Onygenales* and *Eurotiales* are both part of the subclass *Eurotiomycetidae*, this indicates that strong adaptation of the regulation of pentose utilisation has occurred at this evolutionary node. In *Eurotiales* a unique two-component regulatory system for pentose release and metabolism has evolved, while the regulatory system was lost in the *Onygenales*. The observed evolutionary changes (in *Eurotiomycetidae*) mainly affect the regulatory system as in contrast, homologues for most genes of the L-arabinose/D-xylose catabolic pathway are present in all the filamentous fungi, irrespective of the presence of XlnR and/or AraR.

## INTRODUCTION

The order *Eurotiales* consists of the families *Trichocomaceae* and *Elaphomycetaceae*. Most species belonging to the *Trichocomaceae* are saprobic filamentous ascomycetes, which in nature grow predominantly in soil or on decaying plant material. The *Elaphomycetaceae* entails a family of underground, saprobic or mycorrhiza-forming fungi. The family *Trichocomaceae* includes the well-known genera of *Penicillium* and *Aspergillus*. Aspergilli are found throughout the world in almost all ecosystems and are well-known for their ability to degrade different complex plant polymers. Despite the fact that some *Aspergillus* species have evolved additional lifestyles, for example as human or plant pathogens, there seems to be no restriction to a specific niche concerning their saprobic lifestyle.

Decaying plant material consists for a major part of plant cell wall polysaccharides which can be split into three major groups: cellulose, hemicellulose and pectin. L-arabinose and/or D-xylose are the main components of the hemicelluloses arabinoxylan and xyloglucan, and of pectin. Release of these sugars from polysaccharides as well as metabolic conversion of them through the pentose catabolic pathway (PCP) has been studied for many years, particularly in *Aspergillus* and the genus *Trichoderma* belonging to the order *Hypocreales* [reviewed in (de Vries & Visser 2001, de Vries 2003, Stricker *et al.* 2008)]. The PCP was first described in *Aspergillus niger* (Witteveen *et al.* 1989) and shown to consist of a series of reversible reductase/dehydrogenase steps followed by phosphorylation to D-xylulose-5-phosphate, which enters the pentose phosphate pathway (PPP). In *A. niger*, the gene encoding D-xylose reductase (*xyrA*) (Hasper *et al.* 2000), D-xylulokinase (*xkiA*) (van Kuyk *et al.* 2001), L-arabitol dehydrogenase (*ladA*) and xylitol dehydrogenase (*xdhA*) (Seiboth *et al.* 2003, de Groot *et al.* 2007) have been characterised. For *Trichoderma reesei*, genes encoding L-arabitol dehydrogenase (*lad1*) (Richard *et al.* 2001) and xylitol dehydrogenase (*xdh1*) (Seiboth *et al.*

2003) have been described. In *A. niger*, induction of pentose release and the PCP occurs in the presence of L-arabinose and/or D-xylose (Witteveen *et al.* 1989). In the presence of D-xylose, the xylanolytic transcriptional activator XlnR (van Peij *et al.* 1998b) regulates the expression of genes encoding extracellular polysaccharide degrading enzymes, as well as the expression of *xyrA* [reviewed in (de Vries 2003)]. L-arabinose induction of the PCP is not mediated via XlnR. The genes of the L-arabinose catabolic pathway are co-regulated with the genes encoding extracellular arabinolytic enzymes ( $\alpha$ -L-arabinofuranosidase and endoarabinanase) (Flipphi *et al.* 1994, de Vries *et al.* 1994) and L-arabitol is most likely the inducer (de Vries *et al.* 1994, van Kuyk *et al.* 2001). Analysis of *A. niger* arabinolytic regulatory mutants, *araA* and *araB*, demonstrated an antagonistic effect between XlnR and the L-arabinose/L-arabitol responsive regulation (de Groot *et al.* 2003).

In this study, we report the identification and characterisation of the L-arabinose catabolic pathway specific regulator (AraR) in *A. niger* and demonstrate that this regulator is only present in the order *Eurotiales*. These fungi have evolved a fine-tuned two-regulator activating system for pentose release and catabolism compared to other filamentous ascomycetes that only contain XlnR or have neither of the regulators.

## MATERIALS AND METHODS

### Strains, media and growth conditions

The *A. niger* strains used in this study are listed in Table 1 and are all derived from *A. niger* CBS 120.49. *Aspergillus niger* strains were grown in Minimal Medium (MM) or Complete Medium (CM) with addition of a carbon source at 30 °C. MM contained (per liter): 6

**Table 1.** Strains used in this study.

| Strain     | Genotype   | Reference                    |
|------------|--|------------------------------|
| N402       | <i>cspA1</i>   | Bos <i>et al.</i> (1988)     |
| N572       | <i>cspA1, fwnA1, pyrA6, xkiA1, nicA1</i>   | vanKuyk <i>et al.</i> (2001) |
| NW249      | <i>cspA1, ΔargB, pyrA6, nicA1, leuA1</i>   | Jalving <i>et al.</i> (2000) |
| UU-A049.1  | <i>cspA1, pyrA6, nicA1, leuA1, ΔargB::pIM2101(argB+)</i>                               | This study                   |
| UU-A033.21 | <i>cspA1, pyrA6, nicA1, leuA1, ΔargB::pIM2101(argB+), ΔaraR</i>                        | This study                   |
| UU-A054.4  | <i>cspA1, pyrA6, nicA1, leuA1, ΔargB::pIM2101(argB+), ΔaraR::araR</i>                  | This study                   |
| UU-A062.10 | <i>cspA1, ΔargB, nicA1, leuA, pyrA6::A. oryzae pyrA, ΔxlnR</i>                         | This study                   |
| UU-A063.22 | <i>cspA1, nicA1, leuA1, ΔargB::pIM2101(argB+), ΔaraR, pyrA6::A. oryzae pyrA, ΔxlnR</i> | This study                   |

g NaNO<sub>3</sub>, 1.5 g KH<sub>2</sub>PO<sub>4</sub>, 0.5 g KCL, 0.5 g MgSO<sub>4</sub>·7 H<sub>2</sub>O and 200 µl trace elements solution (Vishniac & Santer 1957), pH 6.0. CM = MM supplemented with (L): 2 g peptone, 1 g casamino acids, 1 g yeast extract and 0.5 g yeast ribonucleic acids, pH 6.0. For growth on solid media, 1.5 % agar was added to the medium. When necessary, the medium was supplemented with 0.2 g/L arginine, 0.2 g/L leucine, 0.2 g/L uridine and/or 1 mg/L nicotinamide.

In transfer experiments, all the strains were pre-grown in CM containing 2 % D-fructose. After 16 h of incubation, the mycelium was harvested without suction over a filter, washed twice with MM without a carbon source and transferred to 50 mL MM containing the appropriate carbon source and supplements. The mycelium was harvested with suction over a filter and culture samples were taken after 2 and 4 h of incubation. The mycelium samples were dried between tissue paper and directly frozen in liquid nitrogen.

## Molecular biology methods

Molecular biology methods were performed according to standard procedures (Sambrook *et al.* 1989), unless stated otherwise. All PCR reactions were performed using Accutag™ LA DNA Polymerase (Sigma-Aldrich) according to the manufacturer's instruction. The flanking regions of the *araR* gene were amplified with 5'-primers and 3'-primers (see online Supplemental Table 1) by PCR to generate the 5' flank with the *HindIII/SphI* site and 3' flank with a *KpnI/BamHI* site, respectively, to enable deletion of the complete coding region of *araR* by replacing it with the *argB* selection marker. The functional construct was obtained using PCR with the extreme 5'- end 3'-primers (see online Supplemental Table 1) for complementation of *araR*. The *araR* disruption cassette (containing the *argB* gene for selection for arginine prototrophy) was transformed to the *A. niger* strain NW249 (*pyrA6, leuA1, nicA1, ΔargB*). The *xlnR* gene was amplified with the extreme 5'-primer and 3'-primer by PCR (see online Supplemental Table 1) The PCR fragment was ligated into pGEM-T-easy (Promega) from which the *NsiI/PstI* restriction sites were removed. The construct was digested with *Sall/EcoRI* to remove most of the coding region including the DNA binding domain and ligated with the *A. oryzae pyrA* gene that was digested with *BamHI* (made blunt with Klenow fragment) and *Sall*. The *xlnR* disruption cassette was transformed to *A. niger* strains NW249 (*pyrA6, leuA1, nicA1, ΔargB*) and UU-A033.21 (*pyrA6, leuA1, nicA1, ΔaraR*). All *A. niger* transformations were carried out as described previously (Kusters-van Someren *et al.* 1991).

The primers used to generate the probes for Southern and Northern analysis are listed in online Supplemental Table 1. The

probes were DIG-labelled using the PCR DIG Probe Syntheses Kit (Roche Applied Science) according to the supplier's instructions. A cDNA library (de Groot *et al.* 2007) or genomic DNA (obtained from N402) was used as a template in the PCR reactions for synthesis of the probes.

## Expression analysis

Total RNA was isolated from mycelium that was ground in a microdismembrator (B Braun) using a standard RNA isolation method with the TRIzol Reagent (Invitrogen). In the Northern analysis, 3 µg total RNA was transferred to a Hybond-N<sup>+</sup> membrane (Amersham Biosciences). The Minifold II slot blot apparatus (Schleicher & Schuell) was used for Slot blot analysis. Equal loading was determined by soaking the blot for 5 min in 0.04 % methylene blue, 0.5 M acetate pH 5.2 solution.

Hybridisation of the DIG-labeled probes to the blot was performed according to the DIG user's manual (www.roche-applied-science.com). All the blots were incubated overnight at 50 °C. The blots were exposed for 25 min up to 24 h to a Lumi-Film Chemiluminescent Detection Film (Roche Applied Science). Micro array analysis was performed as described previously (Levin *et al.* 2007).


## Phylogenetic analysis

The amino acid sequences of AraR, XlnR, LadA, XdhA, XyrA and XkiA were used as queries in a local Blast against the protein files of 38 fungal genomes (see online Supplemental Table 2) with a expect value cut-off of 1E-10. The resulting ORFs were aligned using ClustalX and a Maximum Parsimony tree (1 000 bootstraps) was produced using MEGA (v. 4.0).

## Enzyme assays

Extracellular enzyme activity was measured using 0.01 % *p*-nitrophenol linked substrates, 10 µL of the culture samples, 25 mM sodium acetate pH 5.0 in a total volume of 100 µL. Samples were incubated in microtiter plates for 120 min at 30 °C. Reactions were stopped by addition of 100 µL 0.25 M Na<sub>2</sub>CO<sub>3</sub>. Absorbance was measured at 405 nm in a microtiter platereader (Biorad Model 550). The extracellular enzyme activity was calculated using a standard curve ranging from 0 to 80 nmol *p*-nitrophenol per assay volume.

To measure intracellular enzyme activity, cell free extract was prepared by adding 1 mL extraction buffer (50 mM K<sub>2</sub>HPO<sub>4</sub>, 5 mM



**Fig. 1. A.** Expression analysis of the three XlnR homologues (An04g08600 (*araR*), An11g00140 and An11g06290) on D-fructose (1), L-arabinose (2), L-arabitol (3), D-xylose (4) and xylitol (5). **B.** Growth of the reference (UU-A049.1),  $\Delta$ *araR* (UU-A033.21) and  $\Delta$ *araR::araR* (UU-A054.4) on D-glucose and L-arabitol.

MgCl<sub>2</sub>, 5 mM 2-mercaptoethanol, 0.5 mM EDTA) to powdered mycelium. The mixtures were centrifuged for 10 min at 12000 RPM at 4 °C. The L-arabitol and xylitol dehydrogenase activities were determined using 100 mM glycine pH 9.6, 0.4 mM NAD<sup>+</sup> and 1 M L-arabitol or xylitol, respectively. L-arabinose reductase and D-xylose reductase activities were determined using 50 mM Tris-HCl pH 7.8, 0.2 mM NADPH and 1 M L-arabinose or D-xylose, respectively. L-arabinose reductase (ArdA) and D-xylose reductase (XyrA) both convert D-xylose to xylitol and L-arabinose to L-arabitol, but have a higher activity on their primary substrate (de Groot *et al.* 2003). As a result, the measured activity is the sum of the two enzymes. To be able to discriminate between the two enzymes, the ratio of the activity on L-arabinose and on D-xylose was calculated that allows us to extrapolate the relative activities of ArdA and XyrA. An increase in the ratio indicates a relative increase in ArdA or decrease in XyrA, while a reduction in the ratio indicates a relative increase in XyrA or decrease in ArdA.

Absorbance changes were measured at 340 nm using a spectrometer (Spectronic Unicam UV1). L-arabinose and D-xylose reductase activity and L-arabitol and D-xylitol dehydrogenase activity was calculated using the molar coefficient for NADPH and NADH (both  $\epsilon = 6.22 \text{ mM}^{-1}\text{cm}^{-1}$ ) and the following formula:

$$\text{Activity (U/mL)} = [(A/\text{min} - \text{Abl}/\text{min}) * d * v] / (l * a * \epsilon).$$

Abl/min = decrease absorbance per minute before adding substrate.  
A/min = decrease absorbance per minute after adding substrate.  
a = sample volume (mL). d = sample dilution. v = total volume cuvet. l = lightpath (cm). Protein concentrations of intracellular and extracellular samples were determined using a BCA protein assay kit (Pierce).

## RESULTS

### Identification and analysis of *araR*

Blast analysis of XlnR against the *A. niger* genome (Pel *et al.* 2007) revealed 3 homologues with expect values smaller than  $e^{-30}$  (An04g08600, An11g00140, An11g06290). Expression analysis of these genes revealed that the closest *xlnR* homologue (An04g08600) was specifically induced in the presence of L-arabinose or L-arabitol, while only low constitutive expression was observed for An11g06290 and no expression for An11g00140 (Fig. 1A). In order to study its possible role in L-arabinose utilisation,


a disruption strain for An04g08600 (referred to as *araR*) was constructed and verified by Southern analysis (data not shown). The disruption strain showed poor growth on L-arabitol, whereas complementation with *araR* restored growth again (Fig. 1B).

The *araR* gene consists of 2552 bp interrupted by a single intron of 53 bp. Within the 1000 bp promoter region of *araR* putative six binding sites for the carbon catabolite repressor protein CreA (Kulmburg *et al.* 1993) and two binding sites for the xylanolytic regulator XlnR (van Peij *et al.* 1998b, de Vries *et al.* 2002) can be found. The AraR protein contains a Zn(2)Cys(6) binuclear cluster domain (amino acids 36-73, Pfam00172) and a Fungal specific transcription factor domain (amino acids 386-532, Pfam04082). An amino acid motif Arg-Arg-Thr-Leu-Trp-Trp is found at position 493 to 498. This motif differs in only one amino acid from a conserved motif of unknown function found in Zn(2)Cys(6) family members (Arg-Arg-Arg-Leu-Trp-Trp), first described in the UaY regulator in *Aspergillus nidulans* (Suarez *et al.* 1995). AraR shows 32 % identity to XlnR, with the highest homology in the C-terminal part of the proteins. The sequence between the 2<sup>nd</sup> and the 3<sup>rd</sup> Cysteine in the Zn(2)Cys(6) region was previously shown to be important in DNA binding specificity of this class of regulators (Marmorstein *et al.* 1992, Marmorstein & Harrison 1994), but differs significantly between AraR (C<sub>2</sub>HSRRVRC<sub>3</sub>) and XlnR (C<sub>2</sub>NQLRTKC<sub>3</sub>). Between the third and the fourth Cysteine, the Proline residue can be found that is essential for correct folding of the DNA binding domain (Marmorstein *et al.* 1992) and is highly conserved in all the fungal zinc binuclear transcriptional regulators.

### The presence of AraR in the genome is restricted to *Eurotiales* and possibly to *Trichocomaceae*

BlastP analysis of both AraR and XlnR against 38 fungal genome sequences (see online Supplemental Table 2) identified homologues for both proteins in all 11 analysed species of the family *Trichocomaceae* of the order *Eurotiales* (*Aspergillus clavatus*, *A. flavus*, *A. fumigatus*, *A. nidulans*, *A. niger*, *A. oryzae*, *A. terreus*, *Neosartorya fischeri*, *Penicillium chrysogenum*, *P. marneffei*, *Talaromyces stipitatus*), but neither of them was found in three representatives of *Onygenales* (*Coccidioides immitis*, *Histoplasma capsulatum*, *Uncinocarpus reesei*) (Fig. 2). XlnR was also found in the genomes of all other filamentous ascomycetes used in this study. No XlnR and AraR homologues were found in ascomycete yeasts, basidiomycetes or zygomycetes.

In addition, a BlastP analysis was performed with the amino acid sequence of four genes of the *A. niger* pentose catabolic



**Fig. 2.** Bootstrapped (1000 bs) Maximum Parsimony tree of putative homologues of XlnR and AraR in fungi. Homologues of the *A. nidulans* acetate regulatory protein (FacB) were used as an outgroup.



pathway (*ladA*, *xylA*, *xdhA* and *xkiA*) against the genomes of the fungal species that contain XlnR and/or AraR as well as the three *Onygenales* genomes used in this study (see online Supplemental Fig. 1). Phylogenetic analysis showed that all genomes contain homologues of three genes of the pentose catabolic pathway. Homologues for the 4<sup>th</sup> gene (*ladA*) were found in all species except for *Onygenales*.

### Influence of AraR and XlnR on growth of *A. niger* on monomeric and polymeric carbon sources

In addition to the *araR* disruptant (UU-A033.21), an *xlnR* disruptant (UU-A062.10) and an *araR/xlnR* double disruptant (UU-A063.22) were generated, as described in Materials and Methods. The utilisation of several monomeric and polymeric carbon sources was analysed in all strains (including the reference) to determine the effect of the single disruption of the *araR* gene and the double disruption of *araR* and *xlnR* (Fig. 3). Polymeric sugars containing L-arabinose residues (arabinan, Arabic gum, arabinogalactan and apple pectin) and D-xylose residues (birchwood xylan) were included in the analysis. Guar gum was used as a control; it is a galactomannan and contains no L-arabinose or D-xylose residues.

Disruption of *araR* resulted in reduced growth on L-arabinose, xylitol, arabinan, Arabic gum, arabinogalactan and apple pectin and poor growth on L-arabitol (Fig. 3). Disruption of *xlnR* resulted in reduced growth on birchwood xylan, while growth was unaffected on D-xylose, xylitol and the other carbon sources. Disruption of both regulators resulted in a similar phenotype as disruption of *araR* for L-arabitol, Arabic gum, arabinan and arabinogalactan and a similar phenotype as disruption of *xlnR* for birchwood xylan. In contrast to the single disruptants, no growth was observed on D-xylose for the double disruptant, only residual growth on L-arabitol and L-arabinose, and reduced growth on xylitol.

### AraR and XlnR control L-arabinose and D-xylose release and catabolism

The reference,  $\Delta araR$ ,  $\Delta xlnR$  and  $\Delta araR/\Delta xlnR$  strains were pre-grown in complete medium containing D-fructose. After 16 h of growth, equal amounts of mycelium were transferred for 2 and 4 h to minimal medium containing 25 mM D-fructose, 25 mM L-arabinose or 25 mM D-xylose. Extracellular  $\alpha$ -L-arabinofuranosidase (Abf) and intracellular PCP enzyme activities (Ard, Xyr, Lad, Xdh) were analysed. Activity of  $\alpha$ -L-arabinofuranosidase (Abf), L-arabitol dehydrogenase (Lad) and xylitol dehydrogenase (Xdh) was strongly reduced in the  $\Delta araR$  and  $\Delta araR/\Delta xlnR$  strain compared to the reference strain when grown on L-arabinose (Fig. 4A). On D-xylose, Lad and Xdh activity was reduced in  $\Delta araR$  and  $\Delta araR/\Delta xlnR$ . For L-arabinose reductase (ArdA) and D-xylose reductase (XyrA), the ratio of the activity on L-arabinose and on D-xylose was calculated that allowed extrapolation of the relative activities of ArdA and XyrA (see Materials and Methods). The ratio in the  $\Delta araR$  strain became less than 1.0 after 4 h growth in the presence of L-arabinose, while the ratio of the reference strain was around 1.5, which suggests that the ArdA activity was reduced in the  $\Delta araR$  strain (Fig. 4A). The Ard/Xyr ratio in the wild type and  $\Delta araR$  disruptant grown on D-xylose were both around 1. In the absence of both regulators, no Ard and Xyr activities were detected (data not shown). Xylitol dehydrogenase activity (Xdh) was reduced in the  $\Delta araR$  strain on L-arabinose and to a lesser extent on D-xylose compared to the reference strain (Fig. 4). All the measured activities

after 2 h of growth on L-arabinose and D-xylose in the  $\Delta xlnR$  are similar to those published previously (de Groot *et al.* 2003). After 4 h, the difference in activity between the reference and  $\Delta xlnR$  is similar to that observed after 2 h of growth, except for Xdh and Abf. Xdh activity in the  $\Delta xlnR$  became similar to that in the reference strain after 4 h on D-xylose, whereas the Abf activity increased at this point. No activity for any of the enzymes was detected during growth of D-fructose.


In addition, expression levels were determined using micro array analysis for genes involved in release (*abfA*, *abfB*) and catabolism (*ladA*, *xdhA*, *xylA*, *xkiA*) of L-arabinose and D-xylose. No gene expression was observed for any of the genes discussed in this section during growth on 25 mM D-fructose (data not shown). Expression profiles of all the genes in Table 2, except for *araR* and *xlnR*, were confirmed by Northern analysis (see online Supplemental Fig. 1). Expression of *araR* and *xlnR* was below detection levels for Northern analysis in these samples. Disruption of *araR* resulted in 74, 6, 10, 2 and 13-fold reduced expression levels of *abfA*, *abfB*, *ladA*, *xdhA* and *xkiA*, respectively, after 2 h of growth on L-arabinose (Table 2). Disruption of *xlnR* did not significantly reduce expression levels of any of the tested genes, except for *xylA* for which expression reduced 2-fold after 2 h of growth on D-xylose. Disruption of *araR* did not affect *xdhA*, *xkiA* and *xylA* expression on D-xylose, while none of the genes were affected on L-arabinose by disruption of *xlnR* (see online Supplemental Fig. 1). None of the tested genes were expressed in the  $\Delta araR/\Delta xlnR$  strain, except for *abfB* (see online Supplemental Fig. 1). Expression of *xlnR* was not affected in the  $\Delta araR$  on L-arabinose, whereas *araR* expression showed a 3-fold increase in the  $\Delta xlnR$  on D-xylose compared to the reference.

## DISCUSSION


Previously, it has been shown that the pentose catabolic pathway is under control of the D-xylose specific transcriptional activator (XlnR) and a second, unidentified L-arabinose specific transcriptional activator regulator (de Groot *et al.* 2007). In this study, we identified the gene encoding the L-arabinose responsive regulator, AraR, and confirmed its role in the release and catabolism of L-arabinose and D-xylose. AraR is a member of the Zn(2)Cys(6) family of transcriptional regulators and a close homologue of the xylanolytic transcriptional activator XlnR from *A. niger*. Functional analysis of AraR and XlnR as described in this study confirm the previously published antagonistic relation of the two regulatory systems involved in pentose catabolism (de Groot *et al.* 2003).

Expression levels of *abfA*, *abfB*, *ladA* as well as the corresponding enzyme activities (Abf and Lad) were strongly reduced in the  $\Delta araR$  strain on L-arabinose, indicating that they are only controlled by AraR. Gene expression levels of *xdhA* and *xkiA* are reduced in the  $\Delta araR$  strain after 2 h of growth on L-arabinose. On D-xylose, *xdhA* expression is up-regulated in the  $\Delta xlnR$  strain compared to the reference strain, which confirms data published previously (de Groot *et al.* 2007). An increase in *xkiA* expression was observed in the  $\Delta xlnR$  strain on L-arabinose. These results indicate that both AraR and XlnR are involved in regulating the expression of *xdhA* and *xkiA*. The stronger effect in the  $\Delta araR$  strain, suggests that AraR has a larger influence on *xdhA* and *xkiA* expression than XlnR.

Expression of the AraR regulated genes on D-xylose and reduction of the expression in the *araR* disruptant can be explained by the presence of a small amount of L-arabinose in the D-xylose



**Fig. 3.** Growth of the reference strain (Ref., UU-A049.1), and the  $\Delta araR$  (UU-A033.21),  $\Delta xlnR$  (UU-A062.10) and  $\Delta araR/\Delta xlnR$  (UU-A063.22) strains on a selection of mono- and polysaccharides. Concentrations of the substrates were 25 mM for D-glucose, D-xylose, L-arabinose, L-arabitol, xylitol and glycerol, and 1 % for birchwood xylan, Arabic gum, guar gum, arabinan, arabinogalactan and apple pectin.



**Fig. 4.** Comparison of intracellular and extracellular enzyme activities in reference and disruption strains. The reference strains (UU-A049.1),  $\Delta araR$  (UU-A033.21),  $\Delta xlnR$  (UU-A062.10) and  $\Delta araR/\Delta xlnR$  (UU-A063.22) were transferred for 2 and 4 h on 25 mM L-arabinose or 25 mM D-xylose. Extracellular  $\alpha$ -L-arabinofuranosidase (Abf), the ratio of intracellular L-arabinose reductase (ArdA) and D-xylose reductase (XyrA) activity, and the intracellular activities of xylitol dehydrogenase (Xdh) and L-arabitol dehydrogenase (Lad). Black bars: L-arabinose, 2 h; grey bars: L-arabinose, 4 h; dashed bars: D-xylose, 2 h; white bars: D-xylose, 4 h.

preparation from SIGMA (R.P. de Vries, unpubl. data). This is supported by a reduction in the expression of these genes on D-xylose at 4 h compared to 2 h.

The discrepancies between some of the expression and activity data can be explained by the substrate specificities of the enzymes. The L-arabinose and D-xylose reductases are both active on both pentoses, so under conditions where both are expressed, the measured activity is the result of the combined activity of the two enzymes. Although xylitol dehydrogenase is (almost) not active on

L-arabitol, the L-arabitol dehydrogenase is active on xylitol (de Groot *et al.* 2007), indicating that the measured xylitol dehydrogenase can also consist of two components depending on the condition used.

Previously, it has been shown that the expression of *xyrA* was only reduced and not absent in the  $\Delta xlnR$  on D-xylose (de Groot *et al.* 2003) and it was suggested that in addition to *XlnR* another unknown inducing factor is involved. Our results confirm this observation. The reason why there is no reduction in growth of the  $\Delta xlnR$  strain on D-xylose can be explained by the fact



Data from our study was combined with the previously reported data on XlnR (van Peij *et al.* 1998a, de Groot *et al.* 2003) to construct a regulatory model for release and utilisation of L-arabinose and D-xylose in the *A. niger* (Fig. 5). This model correlates not only well with the expression profiles of the pentose-related genes but also with the growth comparison of the disruptant strains and the reference. It indicates that XlnR and AraR control distinct sets of genes in response to the presence of D-xylose and L-arabinose, respectively. However, in the absence of one of the regulators the other can partially compensate for this loss. Although the data supporting this model comes from *A. niger*, we postulate that this model applies to all *Eurotiales*, since we have demonstrated in this study that the presence of AraR is conserved among all species of *Eurotiales* studied so far.



**Table 2.** Expression analysis of genes encoding extracellular L-arabinose releasing enzymes and PCP enzymes. *abfA*, *abfB* =  $\alpha$ -L-arabinofuranosidase A and B, *ladA* = L-arabitol dehydrogenase, *xdhA* = xylitol dehydrogenase, *xyrA* = D-xylose reductase, *xkiA* = D-xylulose kinase, *araR* = arabinolytic regulator, *xlnR* = xylanolytic regulator. The expression levels are mean values of duplicate samples. The ratio was calculated of the expression levels of the reference strain and disruption strain.

|             | Reference<br>2 h L-ara | $\Delta$ <i>araR</i><br>2 h L-ara | Ratio<br>ref/ $\Delta$ <i>araR</i> | Reference<br>2 h D-xyl | $\Delta$ <i>xlnR</i><br>2 h D-xyl | Ratio<br>ref/ $\Delta$ <i>xlnR</i> |
|-------------|------------------------|-----------------------------------|------------------------------------|------------------------|-----------------------------------|------------------------------------|
| <i>abfA</i> | 6622 $\pm$ 919         | 89 $\pm$ 9                        | 74.4                               | 5827 $\pm$ 545         | 7578 $\pm$ 748                    | 0.8                                |
| <i>abfB</i> | 4985 $\pm$ 516         | 901 $\pm$ 143                     | 5.5                                | 869 $\pm$ 4            | 2176 $\pm$ 150                    | 0.4                                |
| <i>ladA</i> | 4224 $\pm$ 417         | 414 $\pm$ 12                      | 10.1                               | 2229 $\pm$ 24          | 3482 $\pm$ 8                      | 0.6                                |
| <i>xdhA</i> | 5013 $\pm$ 661         | 2281 $\pm$ 417                    | 2.2                                | 4344 $\pm$ 315         | 6567 $\pm$ 377                    | 0.7                                |
| <i>xyrA</i> | 4808 $\pm$ 641         | 4048 $\pm$ 685                    | 1.2                                | 6248 $\pm$ 587         | 3655 $\pm$ 66                     | 1.7                                |
| <i>xkiA</i> | 2690 $\pm$ 402         | 211 $\pm$ 18                      | 12.7                               | 2588 $\pm$ 34          | 1843 $\pm$ 102                    | 1.4                                |
| <i>araR</i> | 100 $\pm$ 28           | 1 $\pm$ 0                         | 100                                | 20 $\pm$ 1             | 65 $\pm$ 5                        | 0.3                                |
| <i>xlnR</i> | 145 $\pm$ 21           | 185 $\pm$ 41                      | 0.8                                | 157 $\pm$ 9            | 3 $\pm$ 0                         | 52.3                               |

## ACKNOWLEDGEMENTS

We were supported by grants of the Dutch Foundation for Applied Science (STW) UGC5683 and 07063.

## REFERENCES

- Bos CJ, Debets AJM, Swart K, Huybers A, Kobus G, Slakhorst SM (1988). Genetic analysis and the construction of master strains for assignment of genes to six linkage groups in *Aspergillus niger*. *Current Genetics* **14**: 437–443.
- Flipphi MJA, Visser J, Veen P van der, Graaff LH de (1994). Arabinase gene expression in *Aspergillus niger*: indications for co-ordinated gene expression. *Microbiology* **140**: 2673–2682.
- Gielkens MMC, Visser J, Graaff LH de (1997). Arabinoxylan degradation by fungi: Characterisation of the arabinoxylan arabinofuranohydrolase encoding genes from *Aspergillus niger* and *Aspergillus tubingensis*. *Current Genetics* **31**: 22–29.
- Groot MJL de, Vondervoort PJI van de, Vries RP de, vanKuyk PA, Ruijter GJG, Visser J (2003). Isolation and characterization of two specific regulatory *Aspergillus niger* mutants shows antagonistic regulation of arabinan and xylan metabolism. *Microbiology* **149**: 1183–1191.
- Groot MJL de, Dool C van den, Wösten HAB, Levisson M, vanKuyk PA, et al. (2007). Regulation of pentose catabolic pathway genes of *Aspergillus niger*. *Food Technology and Biotechnology* **45**: 134–138.
- Hasper AA, Visser J, Graaff LH de (2000). The *Aspergillus niger* transcriptional activator XlnR, which is involved in the degradation of the polysaccharides xylan and cellulose, also regulates D-xylose reductase gene expression. *Molecular Microbiology* **36**: 193–200.
- Jalving R, Vondervoort PJI van de, Visser J, Schaap PJ (2000). Characterization of the kexin-like maturase of *Aspergillus niger*. *Applied and Environmental Microbiology* **66**: 363–368.
- Kulmburg P, Mathieu M, Dowzer C, Kelly J, Felenbok B (1993). Specific binding sites in the *alcR* and *alcA* promoters of the ethanol regulon for the CreA repressor mediating carbon catabolite repression in *Aspergillus nidulans*. *Molecular Microbiology* **7**: 847–857.
- Kusters-van Someren MA, Harmsen JAM, Kester HCM, Visser J (1991). The structure of the *Aspergillus niger* *peIA* gene and its expression in *Aspergillus niger* and *Aspergillus nidulans*. *Current Genetics* **20**: 293–299.
- Levin AM, Vries RP de, Conesa A, Bekker C de, Talon M, et al. (2007). Spatial differentiation in the vegetative mycelium of *Aspergillus niger*. *Eukaryotic Cell* **6**: 2311–2322.
- Marmorstein R, Carey M, Ptashne M, Harrison SC (1992). DNA recognition by GAL4: structure of a protein-DNA complex. *Nature* **356**: 408–414.
- Marmorstein R, Harrison SC (1994). Crystal structure of a PPR1-DNA complex: DNA recognition by proteins containing a Zn2Cys6 binuclear cluster. *Genes and Development* **8**: 2504–2512.
- Peij N van, Gielkens MMC, Vries RP de, Visser J, Graaff LH de (1998a). The transcriptional activator XlnR regulates both xylanolytic and endoglucanase gene expression in *Aspergillus niger*. *Applied and Environmental Microbiology* **64**: 3615–3619.
- Peij NNME van, Visser J, Graaff LH de (1998b). Isolation and analysis of *xlnR*, encoding a transcriptional activator coordinating xylanolytic expression in *Aspergillus niger*. *Molecular Microbiology* **27**: 131–142.
- Pel HJ, Winde JH de, Archer DB, Dyer PS, Hofmann G, et al. (2007). Genome sequence of *Aspergillus niger* strain CBS 513.88: a versatile cell factory. *Nature Biotechnology* **25**: 221–231.
- Richard P, Londesborough J, Putkonen M, Kalkkinen N (2001). Cloning and expression of a fungal L-arabinitol 4-dehydrogenase gene. *Journal of Biological Chemistry* **276**: 40631–40637.
- Sambrook J, Fritsch EF, Maniatis T (1989). Molecular cloning -a laboratory manual. Cold Spring Harbour Laboratory, Cold Spring Harbour, N.Y.
- Seiboth B, Harlt L, Pail M, Kubicek CP (2003). D-Xylose metabolism in *Hypocrea jecorina*: Loss of the xylitol dehydrogenase step can be partially compensated for by *lad1*-encoded L-arabinitol-4-dehydrogenase. *Eukaryotic Cell* **2**: 867–875.
- Stricker AR, Mach RL, Graaff LH de (2008). Regulation of transcription of cellulases- and hemicellulases-encoding genes in *Aspergillus niger* and *Hypocrea jecorina* (*Trichoderma reesei*). *Applied Microbiology and Biotechnology* **78**: 211–220.
- Suarez T, Queiroz MV de, Oestreicher N, Scazzocchio C (1995). The sequence and binding specificity of UaY, the specific regulator of the purine utilization pathway in *Aspergillus nidulans*, suggest an evolutionary relationship with the PPR1 protein of *Saccharomyces cerevisiae*. *EMBO Journal* **14**: 1453–1467.
- vanKuyk PA, Groot MJL de, Ruijter GJG, Vries RP de, Visser J (2001). The *Aspergillus niger* D-xylulose kinase gene is co-expressed with genes encoding arabinan degrading enzymes and is essential for growth on arabinose and xylose. *European Journal of Biochemistry* **268**: 5414–5423.
- Vishniac W, Santer M (1957). The thiobacilli. *Bacteriology Reviews* **21**: 195–213.
- Vries RP de (2003). Regulation of *Aspergillus* genes encoding plant cell wall polysaccharide degrading enzymes; relevance for industrial production. *Applied Microbiology and Biotechnology* **61**: 10–20.
- Vries RP de, Flipphi MJA, Witteveen CFB, Visser J (1994). Characterisation of an *Aspergillus nidulans* L-arabitol dehydrogenase mutant. *FEMS Microbiology Letters* **123**: 83–90.
- Vries RP de, Vondervoort PJI van de, Hendriks L, Belt M van de, Visser J (2002). Regulation of the  $\alpha$ -glucuronidase encoding gene (*aguA*) from *Aspergillus niger*. *Molecular and General Genetics* **268**: 96–102.
- Vries RP de, Visser J (2001). *Aspergillus* enzymes involved in degradation of plant cell wall polysaccharides. *Microbiology and Molecular Biology Reviews* **65**: 497–522.
- Witteveen CFB, Busink R, Vondervoort P van de, Dijkema C, Swart K, Visser J (1989). L-arabinose and D-xylose catabolism in *Aspergillus niger*. *Journal of General Microbiology* **135**: 2163–2171.